

EGU2020-4708

<https://doi.org/10.5194/egusphere-egu2020-4708>

EGU General Assembly 2020

© Author(s) 2020. This work is distributed under the Creative Commons Attribution 4.0 License.



Shape asymmetry of rogue waves

Anna Kokorina^{1,2} and Alexey Slunyaev^{1,2,3}

¹Institute of Applied Physics RAS, Nizhny Novgorod, Russian Federation (a.sergeeva@appl.sci-nnov.ru)

²Nizhny Novgorod State Technical University, Nizhny Novgorod, Russian Federation

³National Research University-Higher School of Economics, Nizhny Novgorod, Russian Federation

Direct numerical simulations of the directional sea surface gravity waves are carried out within the framework of the primitive potential equations of hydrodynamics using the High Order Spectral Method. The data obtained for conditions of deep water, the JONSWAP spectrum, and various wave intensities are processed and the results are discussed. The statistical and spectral characteristics of the waves evolve over a long period. The particular asymmetry of the profiles of rogue waves is highlighted. We show that besides the conventional crest-to-trough asymmetry of nonlinear Stokes waves, the extreme events are characterized by a specific combination of the troughs adjacent to the large crest, so that the trough behind the crest is typically deeper than the preceding trough. Surprisingly, the extreme wave crest-to-trough asymmetry and the discrimination between the extreme wave troughs exhibit the tendency to grow when the angle spectrum broadens. This effect contradicts the expectation based on the Benjamin – Feir Index that broad-banded waves should behave similar to linear waves, and hence the asymmetries should diminish.

The research is supported by the RSF grant No. 19-12-00253.

A. Kokorina, A. Slunyaev, The effect of wave nonlinearity on the rogue wave lifetimes and shapes. Proc. 14th Int. MEDCOAST Congress on Coastal and Marine Sciences, Engineering, Management and Conservation (Ed. E. Ozhan), Vol. 2, 711-721 (2019).