

Math-Net.Ru

Общероссийский математический портал

О. В. Починка, С. Ю. Галкина, Д. Д. Шубин, Моделирование градиентно-подобных потоков на n -сфере, *Известия вузов. ПНД*, 2019, том 27, выпуск 6, 63–72

DOI: <https://doi.org/10.18500/0869-6632-2019-27-6-63-72>


Использование Общероссийского математического портала Math-Net.Ru подразумевает, что вы прочитали и согласны с пользовательским соглашением

<http://www.mathnet.ru/rus/agreement>

Параметры загрузки:

IP: 193.238.44.127

7 декабря 2021 г., 15:39:48





УДК 517.9+513.8

<https://doi.org/10.18500/0869-6632-2019-27-6-63-72>

Моделирование градиентно-подобных потоков на n -сфере

O. B. Починка, С. Ю. Галкина, Д. Д. Шубин

Национальный исследовательский университет «Высшая школа экономики»

Россия, 603155 Нижний Новгород, Большая Печерская, 25/12

E-mail: olga-pochinka@yandex.ru, svetlana.u.galkina@mail.ru, schub.danil@yandex.ru

Автор для переписки Данила Денисович Шубин, schub.danil@yandex.ru

Поступила в редакцию 5.09.2019, принята к публикации 28.10.2019, опубликована 2.12.2019

Общей идеей качественного изучения динамических систем, со времён работ А. Андronова, Е. Леонтьевич, А. Майера, является возможность описания динамики системы комбинаторным инвариантом. Так, например, М. Пейштот доказал структурно устойчивые потоки на плоскости определяются единственным образом, с точностью до топологической эквивалентности, классом изоморфных ориентированных графов. Многомерные структурно устойчивые потоки не позволяют ввести их классификацию в рамках общего комбинаторного инварианта. Однако для некоторых подклассов таких систем существует возможность достигнуть полного комбинаторного описания их динамики.

В настоящей работе, основанной на результатах С. Пиллогина, А. Пришляка, В. Гринеса, Е. Гуревич и О. Починка, каждое дерево с двухцветной раскраской вершин реализовано как градиентно-подобный поток на n -сфере, $n > 2$ без гетероклинических пересечений. Эта задача решается с помощью соответствующих операций приклеивания так называемых ячеек Черри к потоку-сдвигу. Этот результат не только завершает топологическую классификацию таких потоков, но и позволяет моделировать системы с регулярным поведением. Для таких потоков реализация особенно важна, поскольку они моделируют, например, процессы присоединения в солнечной короне.

Ключевые слова: градиентно-подобный, моделирование.

Образец цитирования: Починка О.В., Галкина С.Ю., Шубин Д.Д. Моделирование градиентно-подобных потоков на n -сфере // Известия вузов. ПНД. 2019. Т. 27, № 6. С. 63–72. <https://doi.org/10.18500/0869-6632-2019-27-6-63-72>

Финансовая поддержка. Доказательство основной теоремы поддержано Российским Научным Фондом (Грант № 17-11-01041), доказательство вспомогательного результата поддержано Центром фундаментальных исследований Национального исследовательского университета Высшая школа экономики в 2019 году.

Modeling of gradient-like flows on n -sphere

O. V. Pochinka, S. Yu. Galkina, D. D. Shubin

National Research University Higher School of Economics
25/12, Bolshaya Pecherskaya St., 603155 Nizhny Novgorod, Russia
E-mail: olga-pochinka@yandex.ru, svetlana.u.galkina@mail.ru, schub.danil@yandex.ru
Correspondence should be addressed to Danila D. Shubin, schub.danil@yandex.ru
Received 5.08.2019, accepted for publication 28.10.2019, published 2.12.2019

A general idea of the qualitative study of dynamical systems, going back to the works by A. Andronov, E. Leontovich, A. Mayer, is a possibility to describe dynamics of a system using combinatorial invariants. So M. Peixoto proved that the structurally stable flows on surfaces are uniquely determined, up to topological equivalence, by the isomorphic class of a directed graph. Multidimensional structurally stable flows does not allow entering their classification into the framework of a general combinatorial invariant. However, for some subclasses of such systems it is possible to achieve the complete combinatorial description of their dynamics.

In the present paper, based on classification results by S. Pilyugin, A. Prishlyak, V. Grines, E. Gurevich, O. Pochinka, any connected bi-color tree implemented as gradient-like flow of n -sphere, $n > 2$ without heteroclinic intersections. This problem is solved using the appropriate gluing operations of the so-called Cherry boxes to the flow-shift. This result not only completes the topological classification for such flows, but also allows to model systems with a regular behavior. For such flows, the implementation is especially important because they model, for example, the reconnection processes in the solar corona.

Key words: gradient-like flow, modeling.

Reference: Pochinka O.V., Galkina S.Yu., Shubin D.D. Modeling of gradient-like flows on n -sphere. *Izvestiya VUZ. Applied Nonlinear Dynamics*, 2019, vol. 27, no. 6, pp. 63–72. <https://doi.org/10.18500/0869-6632-2019-27-6-63-72>

Acknowledgements. The proof of the main theorem is supported by RSF (Grant no. 17-11-01041), the proof of auxiliary results is supported by Basic Research Program at the National Research University Higher School of Economics (HSE) in 2019.

1. Introduction and statement of results

A general idea of a qualitative study of dynamical systems, going back to the works by A. Andronov [1], E. Leontovich, A. Mayer [2, 3], is a possibility to describe dynamics of a system using combinatorial invariants. A brilliant example of the implementation of such approach is the topological classification of Morse–Smale flows on surfaces obtained by M. Peixoto [4]. He proved that structurally stable flows on surfaces are uniquely determined, up to topological equivalence, by the isomorphic class of a directed graph.

Multidimensional structurally stable flows do not allow to classify them into the framework of a general combinatorial invariant. However, for some subclasses of such systems it is possible to achieve a completely combinatorial description of their dynamics. Thus, according to the results by S. Pilyugin [5], A. Prishlyak [6], V. Grines, E. Gurevich, O. Pochinka [7], the topological equivalence class of a *gradient-like flow* (Morse–Smale flow without periodic orbits) on n -sphere, $n > 2$, without heteroclinic intersections is completely determined by a bi-color tree corresponding to a skeleton of co-dimensional one saddle invariant manifolds. A problem of the realization of an abstract invariant is an integral part of the topological classification. For such flows, it is especially important because they model regular processes in various natural sciences (see, for example, [8]). In particular, such flows model reconnection processes in the solar corona (see, for example, [9, 10]).

In such processes the corona of the Sun is divided on domains by fans and spines (2- and 1-dimensional invariant manifolds) of null points of the magnetic field (the points, at which the strength of magnetic field is null). Restructuring of this domains underlie such effects as the flares and prominences. This energy processes are very important for explanation of many nature laws. The topological structure of a magnetic field is defined by null points, spines, fans and separators, the union of those forming the so-called skeleton of the magnetic field. Experiments and observations show that the evolution of the structure of the magnetic field is similar to relaxation processes. At first plasma evolves slowly for some considerable time but at some point there occurs a topological restructuring of the magnetic configuration (reconnection) [11].

Indeed below we model different stable states of the magnetic field.

Let us recall that *n-ball* or *n-disk* is a manifold with a boundary homeomorphic to a *standard n-ball*

$$\mathbb{B}^n = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_1^2 + \dots + x_n^2 \leq 1\}.$$

An *open n-ball* or *n-disk* we call a manifold homeomorphic to an interior of \mathbb{B}^n . We call by *(n-1)-sphere* a manifold S^{n-1} homeomorphic to a *standard (n-1)-sphere*

$$S^{n-1} = \partial \mathbb{B}^n.$$

Let us consider a class G of gradient-like flows without heteroclinic intersections on n -dimensional sphere S^n , $n \geq 3$, that is, flows whose the non-wandering set consists of a finite many hyperbolic fixed points such that the invariant manifolds of saddle points have no intersections.

Let $f^t \in G$. According to the result by S. Pilyugin [5, Lemma 2.2], the dimensions of the invariant saddle manifolds of f^t have to be only $(n-1)$ and 1. Let us denote by Ω_{f^t} the non-wandering set of f^t , and let

$$\Omega_{f^t}^i = \{p \in \Omega_{f^t} \mid \dim W_p^u = i\}.$$

By [12, Theorem 2.3],

$$S^n = \bigcup_{p \in \Omega_{f^t}} W_p^u = \bigcup_{p \in \Omega_{f^t}} W_p^s.$$

It follows from [13, Proposition 2.3] that for any saddle point σ of a flow f^t the closure of its invariant manifold W_σ^δ with dimension $(n-1)$ contains, except the manifold itself, exactly one fixed point. That point is a sink if $\delta = u$ and a source if $\delta = s$. Then the set $cl W_\sigma^\delta$ is a sphere with dimension $(n-1)$. By [14] and [15] this sphere is cylindrically embedded¹. Denote by m_{f^t} the number of saddle points of a flow f^t . Then the union

$$\mathcal{W}_{f^t} = \bigcup_{p \in \Omega_{f^t}^1} cl W_p^s \cup \bigcup_{q \in \Omega_{f^t}^{n-1}} cl W_q^u$$

of closures of all invariant manifolds of dimension $(n-1)$ divides a sphere S^n into $k_{f^t} = m_{f^t} + 1$ connected components. Denote such components by $D_1, \dots, D_{k_{f^t}}$, and let

$$\mathcal{D}_{f^t} = \bigcup_{i=1}^{k_{f^t}} D_i.$$

¹A sphere $S^{n-1} \subset M^n$ is called *cylindrically embedded* in M^n , if there exists a topological embedding $h : \mathbb{S}^{n-1} \times [-1; +1] \rightarrow M^n$, such that $h(\mathbb{S}^{n-1} \times \{0\}) = S^{n-1}$.




Fig. 1. Example of a flow and its bi-color graph

Рис. 1. Пример потока и его двухцветного графа

A bi-colour graph of a flow $f^t \in G$ is a graph Γ_{f^t} , such that:

- 1) the set $\Gamma_{f^t}^0$ of vertices of Γ_{f^t} bijectively corresponds to \mathcal{D}_{f^t} by a bijection

$$\xi_0: \Gamma_{f^t}^0 \rightarrow \mathcal{D}_{f^t};$$

- 2) two vertices v_i, v_j are connected by an edge $e_{i,j}$ iff domains $D_i = \xi_0(v_i), D_j = \xi_0(v_j)$ have a common boundary;
- 3) an edge $e_{i,j}$ has a colour u (resp. s) if the common boundary of D_i and D_j is the closure of an unstable (resp. stable) saddle manifold (see Fig. 1).

Two graphs Γ_{f^t} and $\Gamma_{f'^t}$ of some flows f^t, f'^t are called *isomorphic* if there exists an isomorphism $\eta: \Gamma_{f^t} \rightarrow \Gamma_{f'^t}$ mapping vertices of Γ_{f^t} into vertices of $\Gamma_{f'^t}$ preserving adjacency and coloring.

It follows from [7], that the flows $f^t, f'^t \in G$ are topologically equivalent iff their graphs Γ_{f^t} and $\Gamma_{f'^t}$ are isomorphic. Indeed, for any flow $f^t \in G$ its bi-color graph is a tree, i.e. connected graph without cycles.

The main result of the present paper is the following theorem.

Theorem 1. *For every bi-color tree Γ there is a flow $f^t \in G$ whose graph Γ_{f^t} is isomorphic to graph Γ .*

Notice that flows of the considered class, under the assumption that they have a unique sink, were classified and realized in [16] by means of a directed graph.

2. Realization of a flow by a bi-color tree

2.1. Description of bi-color tree. Recall some definitions from the graph theory (see, for example, [17] for details).

A *graph* is a pair (V, E) , where V is a set of *vertices* and E is a set of *pairs of vertices*, which are called *edges*. If E contains ordered pairs, then the graph is called a *directed* one. A *k-edge-colouring* of a graph is an assignment of k colours to its edges.

Two vertices are called *adjacent* if they are *connected* by an edge (i.e. they constitute the edge), and the edge is *incident* to each of the vertices. A *loop* is an edge, whose end-vertices coincide. A *simple graph* is an undirected graph without loops.

The number of edges incident to a vertex is called *degree* of the vertex.

A set $\{v_1, (v_1, v_2), v_2, \dots, v_{k-1}, (v_{k-1}, v_k), v_k\}$ is called a *path* of the length k . A path is called a *cycle* if $v_1 = v_k$. A graph is called *connected* if every two its vertices are joined by a path.

A *tree* is a connected acyclic graph. It means that any two its vertices are connected by exactly one path.

Any tree with at least 2 vertices has at least two *pendant vertex*, that is a vertex of the degree 1.

Any tree becomes a *out-tree* if arbitrary its vertex r is selected, as a *root*. In the other words a planted tree is a tree in which one vertex r has been designated as the root and every edge is directed away from the root.

If v is a vertex in a planted tree other than the root, the *parent* of v is the unique vertex w such that there is a directed edge (w, v) . If w is the parent of v , then v is called a *child* of w .

The rooted vertex r by definition has a level 0. The *level d* of any other vertex v in a such planted tree is the number of edges in the unique path between the vertex v and the root r . The *depth* of a tree D is the maximum level of any vertex there.

An *ordered out-tree* is an out-tree where the children of each vertex are ordered.

2.2. Construction of the flow by the graph. To construct a required flow on the n -sphere \mathbb{S}^n for the given bi-color tree Γ choose a pendant vertex r of Γ as a root and an order all children to get from the tree Γ an ordered out-tree. Denote by N the number of all vertices of Γ .

To realize of a flow from the bi-color tree Γ , we will use the idea of embedding of $N - 1$ pairwise disjoint *Cherry boxes* B_v in a *flow-shift* $g_0^t : \mathbb{R}^n \rightarrow \mathbb{R}^n$, given by the formula

$$g_0^t(x_1, \dots, x_n) = (x_1 + t, \dots, x_n)$$

and the cherry box B_v has a form

$$B_v = \{(x_1, \dots, x_n) \in \mathbb{R}^n : |x_1 - \alpha_v| \leq \delta_v, (x_2 - \beta_v)^2 + x_3^2 + \dots + x_n^2 \leq \delta_v^2\}$$

for some $\alpha_v, \beta_v \in \mathbb{R}, \delta_v > 0$ which depends on parameters of v . The dynamics in B_v coincides with the flow-shift dynamics on the boundary of B_v and differs from one inside the box due to the appearance of a saddle and a node. We will say that the dynamics in B_v has a type u (s), if the saddle point has $(n - 1)$ -dimensional unstable (stable) manifold and the node point is a source (sink) (see Fig. 2).

Below we give formulas for the following things:

- 1) The calculation of a position and a size of the Cherry-box B_v ;
- 2) The definition of a flow g_v^t in B_v ;
- 3) The embedding of a resulting dynamics in \mathbb{S}^n .

1. The calculation of a position and a size of the Cherry-box B_v . For the vertex y which is a unique child of the root r we put

$$\alpha_y = 2\rho_y \left(\frac{1}{2} + \frac{1}{2N-4} + \dots + \frac{1}{(2N-4)^{D-2}} \right), \quad \beta_y = 0, \quad \delta_y = 1,$$




Fig. 2. An embedding of Cherry-boxes of the types u and s to the flow-shift

Рис. 2. Вложение ячеек Черри типа u и s в поток-сдвиг

where ρ_y equals 1 (-1) if the edge (r, y) has a colour s (u) and D is the depth of the tree Γ . For any other vertex v with the level $d_v \geq 2$ the parameters of the box B_v are determined through the parameters $\alpha_w, \beta_w, \delta_w$ of its parent's box B_w , the order k_v of v as a child and a number ρ_v which equals 1 (-1) if the edge (w, v) has a colour s (u) by the following way:

$$\delta_v = \frac{\delta_w}{2N-4}, \quad \alpha_v = \rho_v (|\alpha_w| - \delta_w - \delta_v), \quad \beta_v = \beta_w + \frac{\delta_w}{2} - (2k_v - 1)\delta_v.$$

So, the size and position are defined for each Cherry-box corresponding to every vertex of Γ except the root.

2. The definition of a flow g_v^t in B_v . Let

$$\Sigma_v = (x_1 - \alpha_v)^2 + (x_2 - \beta_v)^2 + x_3^2 + \cdots + x_n^2.$$

Define the flow $g_v^t : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by the formulas:

$$\begin{cases} \dot{x}_1 = \begin{cases} 1 - \frac{16\delta_v^2}{9} (\Sigma_v - \delta_v^2)^2, & \Sigma_v \leq \delta_v^2 \\ 1, & \text{otherwise} \end{cases} \\ \dot{x}_2 = \begin{cases} \frac{x_2 - \beta_v}{2} \left(\sin \left(\frac{\pi}{2} \left(\frac{4\Sigma_v}{\delta_v^2} - 3 \right) \right) - 1 \right), & \frac{\delta_v^2}{2} < \Sigma_v \leq \delta_v^2 \\ -(x_2 - \beta_v), & \Sigma_v \leq \frac{\delta_v^2}{2} \\ 0, & \text{otherwise} \end{cases} \\ \dots \\ \dot{x}_n = \begin{cases} \frac{x_n}{2} \left(\sin \left(\frac{\pi}{2} \left(\frac{4\Sigma_v}{\delta_v^2} - 3 \right) \right) - 1 \right), & \frac{\delta_v^2}{2} < \Sigma_v \leq \delta_v^2 \\ -x_n, & \Sigma_v \leq \frac{\delta_v^2}{2} \\ 0, & \text{otherwise} \end{cases} \end{cases}$$

By construction flow g_v^t has exactly two hyperbolic fixed points: the saddle (source) point $P_v(\alpha_v + \rho_v \delta_v / 2, \beta_v, 0, \dots, 0)$ and the sink (saddle) point $Q_v(\alpha_v - \rho_v \delta_v / 2, \beta_v, 0, \dots, 0)$ for $\rho_v = 1$ ($\rho_v = -1$). Define $g_\Gamma^t : \mathbb{R}^n \rightarrow \mathbb{R}^n$ in such a way that it coincides with g_v^t in B_v and is g_0^t outside all Cherry-boxes (see figure 3).




Fig. 3. An example of a tree Γ and the flow g_Γ^t

Рис. 3. Пример дерева Γ и потока g_Γ^t

Let us notice that the flow g_Γ^t has no heteroclinic intersections. Indeed, by the construction the interiors of the Cherry-boxes are pairwise disjoint. Moreover, in the hyperplane $x_1 = \alpha_v$ we have

$$\begin{aligned}\dot{x}_1 < 0 &\quad \text{if } (x_2 - \beta_v)^2 + x_3^2 + \dots + x_n^2 < (\delta_v/2)^2, \\ \dot{x}_1 > 0 &\quad \text{if } (\delta_v/2)^2 < (x_2 - \beta_v)^2 + x_3^2 + \dots + x_n^2 < \delta_v^2.\end{aligned}$$

Also in B_v we have

$$\begin{aligned}\dot{x}_2 \leq 0 &\quad \text{if } x_2 \geq \beta_v, \quad \dot{x}_2 \geq 0 \quad \text{if } x_2 \leq \beta_v, \\ \dot{x}_i \leq 0 &\quad \text{if } x_i \geq 0, \quad \dot{x}_i \geq 0 \quad \text{if } x_i \leq 0, \quad i = 3, \dots, n.\end{aligned}$$

Thus, the invariant $(n-1)$ -manifold of saddle point from B_v outside B_v coincides with a cylinder

$$C_v = \{(x_1, \dots, x_n) \in \mathbb{R}^n : (x_2 - \beta_v)^2 + x_3^2 + \dots + x_n^2 \leq v_v^2\},$$

where $\delta_v/2 < v_v < \delta_v$. By the construction these cylinders are pairwise disjoint, that proves the fact.

3. The embedding of a resulting dynamics in \mathbb{S}^n .

Let us define a flow $h^t : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by the formula:

$$h^t(x_1, x_2, \dots, x_n) = (2^t x_1, 2^t x_2, \dots, 2^t x_n).$$

Let $\mathbb{R}_+^n = \{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 \geq 0\}$ and $C = \{(x_1, \dots, x_n) \in \mathbb{R}^n : x_2^2 + \dots + x_n^2 \leq 1\}$. It is easy to verify that a diffeomorphism $\zeta : \mathbb{R}_+^n \setminus O \rightarrow C$ given by the formula

$$\zeta(x_1, \dots, x_n) = \left(\log_2 \varrho, \frac{x_2}{\varrho}, \dots, \frac{x_n}{\varrho} \right), \quad \varrho = \sqrt{x_1^2 + \dots + x_n^2}$$

conjugates the diffeomorphisms $h^t|_{\partial \mathbb{R}_+^n \setminus O}$ and $g^t|_{\partial C}$. It allows to define a flow $\varphi^t : \mathbb{R}^n \rightarrow \mathbb{R}^n$ in such a way that φ^t coincides with h^t outside $\text{int } \mathbb{R}_+^n$ and coincides with $\zeta^{-1} g_\Gamma^t \zeta$ on \mathbb{R}_+^n .

Let us project the flow φ^t to the n -sphere by means the stereographic projection.

Denote by $\mathcal{N}(0, \dots, 0, 1)$ the North Pole

of the sphere \mathbb{S}^n . For every point $x = (x_1, \dots, x_{n+1})$ in $\mathbb{S}^n \subset \mathbb{R}^{n+1}$ there is the unique line passing through the points \mathcal{N} and x . This line intersects $\mathbb{R}^n = O x_1 \dots x_n$ at exactly one point $\vartheta(x)$ (see figure 4), which is called *the stereographic projection of the point x* . One can easily to check that $\vartheta : \mathbb{S}^n \setminus \{\mathcal{N}\} \rightarrow \mathbb{R}^n$ is a diffeomorphism, given by the formula

$$\vartheta(x_1, \dots, x_{n+1}) = \left(\frac{x_1}{1 - x_{n+1}}, \dots, \frac{x_{n-1}}{1 - x_{n+1}}, \frac{x_n}{1 - x_{n+1}} \right).$$

As flow φ^t coincides with h^t in some neighborhoods of the origin O and of the infinity point, hence, it induces on \mathbb{S}^n the required flow

$$f^t(x) = \begin{cases} \vartheta^{-1}(\varphi^t(\vartheta(x))), & x \neq \mathcal{N}; \\ \mathcal{N}, & x = \mathcal{N}. \end{cases}$$




Fig. 4. The stereographic projection

Рис. 4. Стереографическая проекция

References

1. Andronov A.A. and Pontryagin L.S. Rough systems. *Doklady Akademii nauk SSSR*, 1937, vol. 14, no. 5, pp. 247–250 (in Russian).
2. Leontovich E.A., Mayer A.G. About trajectories determining qualitative structure of sphere partition into trajectories. *Doklady Akademii nauk SSSR*, 1937, vol. 14, no. 5, pp. 251–257 (in Russian).
3. Leontovich E.A., Mayer A.G. About scheme determining topological structure of partition into trajectories. *Doklady Akademii nauk SSSR*, 1955, vol. 103, no. 4, pp. 557–560 (in Russian).
4. Peixoto M. On the Classification of Flows on Two Manifolds. *Dynamical systems Proc.*, 1971.
5. Pilyugin S.Yu. Phase diagrams determining Morse–Smale systems without periodic trajectories on spheres. *Diferencial'nye uravneniya*, 1978, vol. 14, no. 2, pp. 245–254 (in Russian).
6. Prishlyak A.O. Morse–Smale vector-fields without closed trajectories on three-dimensional manifolds. *Matematicheskie zametki*, 2002, vol. 71, no. 2, pp. 254–260 (in Russian).
7. Grines V.Z., Gurevich E.Ya., Pochinka O.V. A combinatorial invariant of Morse–Smale diffeomorphisms without heteroclinic intersections on the sphere S^n , $n \geq 4$. *Math. Notes*, 2019, vol. 105, no. 1, pp. 132–136.
8. Pesin Ya.B., Yurhchenko A.A. Some physical models described by the reaction-diffusion equation and chains of connected maps. *UMN*, 2004, vol. 59, no. 3(357), pp. 81–114 (in Russian).
9. Browns D.S., Priest E.R. The topological behaviour of stable magnetic separators. *Sol. Phys.*, 1999, vol. 190, pp. 25–33.
10. Grines V., Zhuzhoma E.V., Pochinka O., Medvedev T.V. On heteroclinic separators of magnetic fields in electrically conducting fluids. *Physica D: Nonlinear Phenomena*, 2015, vol. 294, pp. 1–5.
11. Priest E., Forbes T. Magnetic Reconnection: MHD Theory and Applications. Cambridge Univ. Press, Cambridge, 2000.
12. Smale S. Differentiable dynamical systems. *Bull. Amer. Soc.*, 1967, vol. 73, pp. 747–817.
13. Grines V.Z., Medvedev T.V., Pochinka O.V. Dynamical Systems on 2- and 3- Manifolds. *Dev. Math.*, 46, Springer, Cham, 2016, xxvi+295 pp.
14. Cantrell J.C. Almost locally flat sphere S^{n-1} in S^n . *Proceeding of the American Mathematical society*, 1964, vol. 15, pp. 574–578.
15. Brown M. Locally Flat Imbeddings of Topological Manifolds. *Annals of Mathematics Second Series*, 1962, vol. 75, pp. 331–341.
16. Grines V.Z., Gurevich E.Ya., Medvedev V.S. Classification of Morse–Smale diffeomorphisms with one-dimensional set of unstable separatrices. *Proc. Steklov Inst. Math.*, 2010, vol. 270, pp. 57–79.
17. Harary F. Graph Theory. Addison-Wesley, Reading, MA, 1969.

Библиографический список

1. Андронов А.А., Понtryагин Л.С. Грубые системы // Доклады академии наук СССР. 1937. Т. 14, № 5. С. 247–250.
2. Леонтович Е.А., Майер А.Г. О траекториях, определяющих качественную структуру разбиения сферы на траектории // ДАН СССР. 1937. Т. 14, № 5. С. 251–257.
3. Леонтович Е.А., Майер А.Г. О схеме, определяющей топологическую структуру разбиения на траектории // ДАН СССР. 1955. Т. 103, № 4. С. 557–560.

4. Peixoto M. On the Classification of Flows on Two Manifolds. *Dynamical systems Proc.*, 1971.
5. Пилигин С.Ю. Фазовые диаграммы, определяющие системы Морса–Смейла без периодических траекторий на сферах // Дифференц. уравнения. 1978. Vol. 14, № 2. С. 245–254.
6. Прииляк А.О. Векторные поля Морса–Смейла без замкнутых траекторий на трехмерных многообразиях // Матем. заметки. 2002. Vol. 71, № 2. С. 230–235.
7. Grines V.Z., Gurevich E.Ya., Pochinka O.V. A combinatorial invariant of Morse–Smale diffeomorphisms without heteroclinic intersections on the sphere S^n , $n \geq 4$ // Math. Notes. 2019. Vol. 105, no. 1. P. 132–136.
8. Песин Я.Б., Юрченко А.А. Некоторые физические модели, описываемые уравнением реакции–диффузии, и цепочки связанных отображений // УМН. 2004. 59:3(357). С. 81–114.
9. Browns D.S., Priest E.R. The topological behaviour of stable magnetic separators // Sol. Phys. 1999. Vol. 190. P. 25–33.
10. Grines V., Zhuzhoma E.V., Pochinka O., Medvedev T.V. On heteroclinic separators of magnetic fields in electrically conducting fluids // Physica D: Nonlinear Phenomena. 2015. Vol. 294. P. 1–5.
11. Priest E., Forbes T. Magnetic Reconnection: MHD Theory and Applications. Cambridge univ. Prees, Cambridge, 2000.
12. Smale S. Differentiable dynamical systems // Bull. Amer. Soc. 1967. Vol. 73. P. 747–817.
13. Grines V.Z., Medvedev T.V., Pochinka O.V. Dynamical Systems on 2- and 3- Manifolds // Dev. Math., 46, Springer, Cham, 2016, xxvi+295 pp.
14. Cantrell J.C. Almost locally flat sphere S^{n-1} in S^n // Proceeding of the American Mathematical society. 1964. Vol. 15. P. 574–578.
15. Brown M. Locally Flat Imbeddings of Topological Manifolds // Annals of Mathematics Second Series. 1962. Vol. 75. P. 331–341.
16. Grines V.Z., Gurevich E.Ya., Medvedev V.S. Classification of Morse–Smale diffeomorphisms with one-dimensional set of unstable separatrices // Proc. Steklov Inst. Math. 2010. Vol. 270. P. 57–79.
17. Harary F. Graph Theory. Addison-Wesley, Reading, MA, 1969.

Починка Ольга Витальевна – родилась в 1972 году. Окончила механико-математический факультет Нижегородского государственного университета им. Н.И. Лобачевского (1994), аспирантуру (НГСХА, 2004) и докторантуру (ННГУ, 2011). Защищила диссертации на соискание ученых степеней кандидата физико-математических наук (НГСХА, 2004) и доктора физико-математических наук (ННГУ, 2012). С 1997 по 2014 год работала на кафедре теории функций механико-математического факультета ННГУ им. Н.И. Лобачевского: сначала в должности ассистента, затем старшего преподавателя, доцента. С 2014 года по настоящее время – профессор, заведующая кафедрой фундаментальной математики Национального исследовательского университета Высшая школа экономики. Профессиональные интересы – качественная теория динамических систем.

Россия, 603155 Нижний Новгород, Большая Печерская, 25/12
Лаборатория топологических методов в динамике, Национальный
исследовательский университет «Высшая школа экономики»
E-mail: olga-pochinka@yandex.ru



Галкина Светлана Юрьевна – родилась в 1963 году. Окончила механико-математический факультет Московского государственного университета им. М.В. Ломоносова (1986) и аспирантуру (МГУ, 1991). Кандидат физико-математических наук (МГУ, 2002). С 1991 по 2005 год работала в Нижегородском государственном педагогическом университете, на кафедре математического анализа: сначала в должности ассистента, затем старшего преподавателя, доцента, и.о. заведующего кафедрой математического анализа. С 2005 по 2019 год – доцент кафедры прикладной математики Нижегородского государственного университета им. Н.И. Лобачевского. С 2019 г. по настоящее время – доцент кафедры фундаментальной математики Национального исследовательского университета Высшая школа экономики. Научные интересы: функциональный анализ, теория функций действительного переменного, теория аппроксимации функций.

Россия, 603155 Нижний Новгород, Большая Печерская, 25/12
Национальный исследовательский университет «Высшая школа экономики»
E-mail: svetlana.u.galkina@mail.ru



Шубин Данила Денисович – родился в городе Павлово Нижегородской области (2000). С 2018 г. по настоящее время – студент факультета математики, информатики и компьютерных наук НИУ ВШЭ (Нижний Новгород), стажёр-исследователь лаборатории топологических методов в динамике (ИМиКН НИУ ВШЭ).

Россия, 603155 Нижний Новгород, Большая Печерская, 25/12
Лаборатория топологических методов в динамике, Национальный
исследовательский университет «Высшая школа экономики»
E-mail: schub.danil@yandex.ru

